

Product Instructions.

Title: "FRAN"

Release form: Rectal suppositories.

Composition: One suppository contains:

The active ingredient is *lactoimmunoglobulins* - 100 mg.

Excipients: cocoa butter 2g.

Packing: lacto candles, 10 - 30 pieces per pack.

Pharmacological action:

The drug "FRAN" belongs to the group of targeted, immuno-biological, selectively acting drugs. The active principle of the drug "FRAN" are antibodies that "target" various structures of cell walls (receptors, markers), directly or indirectly associated with the tumor process.

The mechanism of action of targeted drugs:

There are various structures (receptors, transmembrane proteins, etc.) on the surface of cells, necessary for their vital activity. The metabolic processes of tumor cells differ from the metabolism of ordinary cells, which causes the formation on their membranes of a large number of biological structures associated with tumor metabolic processes.

Produced antibodies selectively interact with pathological structures (receptors, markers) of the cell.

The number of pathological structures on the cell surface determines the density of antibodies bound to them. At a high density of antibodies, the subunits of the natural component of blood serum get activated and assembled - a complement that destroys the cell wall and causes its death (lysis).

Indications:

- Benign and malignant tumors of various localization, breast cancer and other oncological diseases;
- Autoimmune diseases;
- Multiple sclerosis;
- Diabetes;
- Allergic diseases;
- Erosion of the cervix, herpes;

- Parkinson's disease;
- Alzheimer's disease;
- Rheumatism, arthritis, arthrosis;
- Osteochondrosis;
- Hepatitis C;
- Bronchial asthma;
- Prostatitis;
- Dystrophy.

Contraindications:

No allergic reactions were detected during the tests.

Method of application:

Suppositories "FRAN" are inserted rectally after bowel movement. Before inserting a candle, it is recommended to freeze it in the freezer compartment of the refrigerator to prevent melting from hand heat.

After the insertion of the suppository, it is recommended to lie down on the side for 20 - 30 minutes, for better absorption of the suppository components in the intestines.

A break in the use of the drug is allowed for no more than 3 days. In case of a break, use a double dose at once (2 suppositories at a time).

In case of a break of more than 3 days, you can continue treatment after 7, 14 or 21 days (in multiples of 7 days).

A repeated course, with the testimony of the attending physician, can be started immediately, without interruption.

It is advisable not to interrupt the course unnecessarily.

Dosage: 1 or 2 suppositories daily, unless otherwise prescribed by the doctor.

Prescription of the drug depends on the severity and development of the disease. The minimum course of treatment is 30 suppositories.

During pregnancy: it is not advisable to use the drug, as research has not been carried out.

When breastfeeding: there are no contraindications for use.

Side effects: a slight increase in temperature and a slight exacerbation of the disease in the first days of usage are possible.

Special instructions:

Attention !!! When undergoing a course of chemotherapy, first the doctor must set the dosage of chemotherapy and only after that the drug "FRAN" can be used, otherwise the wrong dosage of chemotherapy may be set, exceeding the norm by several times.

When treating allergies, it is advisable to cause mild allergic reactions while using the "FRAN" preparation, in order to train the body's immunity.

During the period of use of the drug FRAN, do not cancel the drugs that were prescribed by the attending physician. If you have any questions, contact the seller's consultant.

Interaction with other drugs: It is not recommended to use "FRAN" suppositories in conjunction with other immuno-modulating drugs.

Compatible with other anticancer drugs and chemotherapeutic agents.

Overdose: When using a dose 5 times higher than the recommended one, no side effects have been identified.

Expiration date: 24 months

Storage conditions: store in a dark place, out of reach of children, at a temperature not higher than +25 degrees. Do not expose to heat above + 50 degrees. Prevent direct sunlight.